

Study of the Coleman - de Luccia instanton of the second order

Michal Demetrian*
 Comenius University
 Mlynska Dolina F2, 842 48,
 Bratislava, Slovak Republic
 (Dated: February 7, 2008)

We study the second order Coleman - de Luccia instanton which appears as the curvature of the effective potential reaches a sufficiently large value. We show how one can find the approximative formula for this instanton by perturbative expansion in the case when the second derivative of the effective potential divided by the Hubble parameter squared is close to -10 , and we perform a numerical study of this instanton in the case of quasi-exponential potential.

PACS numbers: 98.80.Cq

I. INTRODUCTION

The theory of false vacuum decay in de Sitter universe via formation of rapidly expanding bubbles was introduced by Coleman and de Luccia [1] and it has been studied intensively as an ingredient of old inflation [2] and of open inflation [3, 4]. This mode of vacuum decay is described by the Coleman - de Luccia (CdL) instanton - the nontrivial finite-action $O(4)$ solution to the Euclidean equations for coupled scalar and gravitational fields, with an effective potential containing a true vacuum (at which it has the global minimum equal to zero) and a false vacuum (at which it has a local minimum) separated from the true vacuum by a finite potential barrier. It is given by two functions $\Phi(\tau)$ and $a(\tau)$, where Φ is the scalar field, a is the radius of the 3-spheres of homogeneity determined from the circumference and τ is the radius of the 3-spheres of homogeneity measured from the centre. Denote the effective potential by V . Functions Φ and a obey the equations

$$\ddot{\Phi} + \frac{3\dot{a}}{a} = V', \quad \ddot{a} = -\frac{8\pi}{3} (\dot{\Phi}^2 + V) a, \quad (1)$$

where the overdot denotes differentiation with respect to τ and the prime differentiation with respect to Φ . The boundary conditions ensuring finiteness of the (Euclidean) action

$$S_E = \int d^4x \sqrt{|g_E|} \left[\frac{1}{2} g_E(\partial\Phi, \partial\Phi) + V(\Phi) - \frac{R(g_E)}{16\pi} \right].$$

are

$$a(0) = \dot{\Phi}(0) = \dot{\Phi}(\tau_f) = 0 \text{ and } \dot{a}(0) = 1, \quad (2)$$

where $\tau_f > 0$ is defined by the equation $a(\tau_f) = 0$.

Being motivated by the arguments from the papers [5, 6, 7, 8] the authors of the paper [9] have proved

both a necessary and a sufficient condition for the existence of the CdL instanton. Before we formulate them let us introduce useful notations which will be used throughout the paper: by $V_M = V(\Phi_M)$ we denote the local maximum of V (the top of the barrier); and by $H(\Phi)$ we mean the value of the Hubble parameter corresponding to the energy density equal to $V(\Phi)$, $H(\Phi) = \sqrt{8\pi V(\Phi)/3}$. The conditions read:

If the inequality $-V''_M/H_M^2 > 4$ holds then the CdL instanton exists.

If the CdL instanton exists then the inequality $-V''(\Phi)/H^2(\Phi) > 4$ holds somewhere in the barrier.

For a given theory (potential) the conditions leave a gray zone in the parameter space of the potential in which they cannot decide whether the CdL instanton exists or not. It has been shown in [9] that in addition to the first order instanton, also instantons of higher order exist if the curvature of the potential in its local maximum is sufficiently large. Namely, any finite-action solution of Eqs. (1) can be characterized by how many times the scalar field crosses Φ_M . If the scalar field crosses Φ_M l times, we shall call the solution the l th order CdL instanton. From the analysis of the instantons localized close to the top of the barrier [9] it follows that

If $-V''_M/H_M^2 > l(l+3)$ then the potential admits the l th order instanton.

For odd l there is, for given V , at most one instanton of l th order; for even l we have two l th order instantons. One of them starts and ends at Φ_i^L less than Φ_M and the other one starts and ends at Φ_i^R greater than Φ_M .

II. ANALYTIC APPROACH TO THE SECOND ORDER CDL INSTANTON LOCALIZED CLOSE TO THE TOP OF THE POTENTIAL BARRIER

The aim of this paragraph is to solve perturbatively Eqs. (1) in the case when $-V''_M/H_M^2 \approx 10$. Our task

*Electronic address: demetrian@fmph.uniba.sk

is to sketch how to get such a solution. The detailed explanation of this method (together with its application to the first order CdL instanton) will appear in [10].

It is useful to introduce the notations

$$\begin{aligned} y(x) &\equiv \Phi(x) - \Phi_M = \sum k^n u_n(x) \\ a(x) &= C H_M^{-1} \sum k^n v_n(x) \\ -\frac{V_M''}{H_M^2} &= 10 + \sum k^n \Delta_n, \end{aligned}$$

where we are using the dimensionless Euclidean time $x = H_M \tau$ that runs from 0 to π , $C = 8\pi/3$, and k is our parameter of perturbative expansion - i.e. the amplitude of $\Phi - \Phi_M$. We will continue our calculations up to the order in which we will be able to decide what quantity seems to be interesting or helpful if one would like to construct (numerically) the second order instanton in a given theory. The functions u_n and v_n obey differential equations

$$\begin{aligned} u_n''(x) + 3 \cot(x) u_n'(x) + 10 u_n(x) &= \mathcal{U}_n(x), \\ v_n''(x) + v_n(x) &= \mathcal{V}_n(x) \sin(x), \end{aligned} \quad (3)$$

where the functions $\mathcal{U}_n$ and $\mathcal{V}_n$ can be computed order by order from the system of equations

$$\begin{aligned} y''(x) + \frac{3a'(x)}{a(x)} y'(x) &= \frac{V'(y)}{H_M^2}, \\ a''(x) &= -C \left(y'^2(x) + \frac{V(y)}{H_M^2} \right) a(x). \end{aligned}$$

Since $\mathcal{U}_1$ is identically equal to zero, u_1 is the nonsingular (at the end points) solution of the equation

$$u_1''(x) + 3 \cot(x) u_1'(x) + 10 u_1(x) = 0$$

i.e.

$$u_1(x) = \frac{1}{4} (5 \cos^2(x) - 1). \quad (4)$$

To get v_2 one has to find out that $\mathcal{V}_2 = -(u_1')^2 + 5u_1^2$ and therefore v_2 obeys

$$v_2''(x) + v_2(x) = \frac{5}{128} \left(-7 \sin(x) + \frac{15}{2} \sin(3x) + \frac{45}{2} \sin(5x) \right)$$

with the initial conditions $v_2(0) = v_2'(0) = 0$. We straightforwardly obtain that

$$\begin{aligned} v_2(x) &= \frac{5}{256} \left(7x \cos(x) - \frac{15}{8} (\sin(3x) + \sin(5x)) \right. \\ &\quad \left. + 8 \sin(x) \right). \end{aligned} \quad (5)$$

We can proceed to u_2 ; in order to do this we need to know the function $\mathcal{U}_2$,

$$\mathcal{U}_2(x) = -\Delta_1 u_1(x) + \frac{1}{2} \eta u_1^2(x),$$

where $\eta = V_M''' / H_M^2$. The equation for u_2 reads

$$\begin{aligned} u_2''(x) + 3 \cot(x) u_2'(x) + 10 u_2(x) &= -\frac{\Delta_1}{4} (5 \cos^2(x) - 1) \\ &\quad + \frac{\eta}{32} (5 \cos^2(x) - 1)^2. \end{aligned}$$

The last equation can be transformed to an equation of hypergeometric type by changing the independent variable $z = \cos(x)$. In this way one gets

$$\begin{aligned} (1 - z^2) u_2''(z) - 4z u_2'(z) + 10 u_2(z) &= \\ -\frac{\Delta_1}{4} (5z^2 - 1) + \frac{\eta}{32} (5z^2 - 1)^2. \end{aligned}$$

We need a solution of this equation which is bounded (at the points $z = \pm 1$). Such a solution is given by a polynomial; one can easily see that it must have the form $\alpha + \beta z^4$. Inserting the previous ansatz in to the equation we get a system of three linear equations for α , β and Δ_1 . These equations are parametrized by η and their solution reads

$$\Delta_1 = \frac{1}{6} \eta \quad (6)$$

and

$$u_2(x) = \frac{\eta}{192} \left(\frac{7}{5} - \frac{25}{3} \cos^4(x) \right). \quad (7)$$

The "quantization rule" for k follows from Eq. (6),

$$k = -\frac{6}{\eta} \left(\frac{V_M''}{H_M^2} + 10 \right). \quad (8)$$

It is also possible, within our perturbative approach, to answer the question how the action of our second order CdL instanton differs from that quantity of related Hawking - Moss instanton. In order to find this difference one has to proceed to the third order calculations and obtain the function v_3 . However, we are not going to do this. Comparison of the two actions is going to appear in [10].

We can conclude that, in addition to the parameter $V_M'' / H_M^2 + 10$, our results depend crucially on the value of the third derivative of the potential at Φ_M . Moreover, we can see from Eq. (8) that these results do not hold in the case $V_M''' = 0$ and they require, for a given $V_M'' / H_M^2 + 10$, a sufficiently large value of η . To determine approximative behaviour of the second order CdL instanton in the case when η is close to zero one should perform extended analysis including the solution of the instanton's equations, at least, up to the third order in k (to determine Δ_2). Instead of doing this, we are going to investigate the properties of the second order CdL instanton numerically, and we understand previously presented steps as a help in numerical calculations. Namely, they showed us that the third derivative of the effective potential in its local maximum can be regarded as an interesting quantity.

III. NUMERICAL STUDY OF THE SECOND ORDER CDL INSTANTON

Let us consider, as an example, the theory with the potential

$$V(\Phi) = \left(V_0 + \frac{1}{2}\Phi^2 \right) e^{-\Phi/\Phi_0}, \quad (9)$$

where V_0 and Φ_0 are positive constants. The motivation for this potential can be found in [11]. The potential (9) is also very convenient for the investigation of the CdL instanton of the second order because it allows us to preserve the barrier while V_M''' is running from negative to positive values; quartic potential has not this property. The barrier in the theory (9) exists only if $V_0 < 1/2\Phi_0^2$, and the values of Φ at which V has the local maximum and local minimum respectively are

$$\Phi_{m,M} = \Phi_0(1 \mp \Delta),$$

where we have introduced the parametrization of the potential by (Φ_0, Δ) with

$$\Delta = \sqrt{1 - 2\frac{V_0}{\Phi_0^2}}, \quad \Delta \in [0, 1].$$




FIG. 1: Dependence of the second order CdL instantons in the theory (9) on the parameter Δ . Description of the quantities introduced in this figure: Φ_i^L is the initial and final value of Φ for the instanton starting left from Φ_M , Φ_{max}^L is the maximal value of Φ for this instanton; Φ_i^R is the initial and final value of Φ for the instanton starting right from Φ_M and Φ_{min}^R is the minimal value of Φ for this instanton. We can see that the right-handed instanton gets shrunk (in the Φ direction) towards Φ_M as Δ approaches 1 and the left-handed instanton is the narrowest as Δ approaches 0. The absolute value of the difference $\Phi_i^L - \Phi_M$ falls down as $1/V_M'''$ for $\Delta \rightarrow 0^+$ and analogically $\Phi_i^R - \Phi_M$ falls down for $\Delta \rightarrow 1^-$. This means that in the case $V_M''' < 0$ ($V_M''' > 0$) the left (right)-handed instanton is the instanton that approaches the limit instanton as $-V_M''/H_M^2 - 10 \rightarrow 0$.

The quantities interesting for our goal are

$$-\frac{V_M''}{H_M^2} = \frac{3\Delta}{8\pi\Phi_0^2(1+\Delta)}$$

and

$$V_M''' = \frac{2\Delta - 1}{\Phi_0^2} \exp(-1 - \Delta).$$

We have put in our numerical computations $-V_M''/H_M^2 - 10 = 0.1$. We investigated parameters of the CdL instantons of the second order as functions of the parameter Δ in its range of definition. Remind that there are always two such instantons - see the introduction. The results are plotted on the fig. 1.

Acknowledgments

The author thanks Vlado Balek for useful discussions. This paper was supported by the grant VEGA 1/3042/06.

-
- [1] Coleman, S. and de Luccia, F., Phys. Rev. **D21**, 3305 (1980).
 - [2] Guth, A.H., Phys. Rev. **D23**, 347 (1981).
 - [3] Bucher, M., Goldhaber, A.S. and Turok, N., Phys. Rev. **D52**, 3314 (1995).
 - [4] Linde, A.D. and Mezhlumian, A., Phys. Rev. **D52**, 6789 (1995).
 - [5] Hawking, S.W. and Moss, I.G., Phys. Lett. **110B**, 35 (1982).
 - [6] Linde, A.D., "Particle physics and inflationary cosmology" (Harwood, Chur, Switzerland 1990).
 - [7] Jensen, L. and Steinhardt, P.J., Nucl. Phys. **B237**, 176 (1984).
 - [8] Goncharov, A.S. and Linde, A.D., Fiz. Elem. Chast. At. Yad. **17**, 837 (1986).
 - [9] Balek, V. and Demetrian, M., Phys.Rev. **D69**, 063518 (2004).
 - [10] Balek, V. and Demetrian, M., Phys.Rev. **D71**, 023512 (2005).
 - [11] Parsons, P. and Barrow, J.D., Phys. Rev. **D51**, 6757 (1995).